



SCREENING OF NUTRITIONAL COMPONENTS FOR α -AMYLASE PRODUCTION IN SUBMERGED FERMENTATION BY BACTERIA ISOLATED FROM SOIL USING PLACKETT-BURMAN DESIGN

SUJATA D. BANSODE

Amrutvahini College of Pharmacy, Sangamner-422 608, India, E. mail: suja_bansode@rediffmail.com

ABSTRACT

Thermostable α -amylase can be produced by the bacteria isolated from soil in a liquid media. Screening of important nutrient parameters from selected seven medium components viz. starch, yeast, extract, tryptone, K_2HPO_4 , $CaCl_2 \cdot 2H_2O$, $MgSO_4 \cdot 7H_2O$ and KCl were carried out in shake flask cultures for enzyme production using Plackett-Burman experimental design of twelve trials. Among these starch, yeast extract and $CaCl_2 \cdot 2H_2O$ contributes to large extent; tryptone and K_2HPO_4 have moderately significant; while $MgSO_4 \cdot 7H_2O$ and KCl shows minute effect for amylase production by submerged fermentation.

Keywords: α -Amylase, submerged fermentation, Plackett-Burman design

INTRODUCTION

Microorganisms had significant contribution in production of various industrial enzymes. The global market for industrial enzymes estimated at \$2 billion in 2004 and expected to rise at an average annual growth rate of 3.3%¹. These starch-degrading amyolytic enzymes are of great significance in biotechnological applications ranging from food, fermentation, textile, paper, pharmaceutical to sugar industries^{2,3,4}. Conversion of starch into sugar, syrups and dextrans forms the major part of starch processing industry⁵. Starch degrading enzymes like amylase have received great deal of attention because of their perceived technological significance and economic benefits⁶. Nowadays, amylases (α -amylases, β -amylases and glucoamylases) represent one of the most important enzyme groups within the field of biotechnology⁷. α -amylase (EC 3.2.1.1, 1,4- α -D-glucan glucohydrolase, endoamylase) is a classical calcium-containing enzyme catalyze hydrolysis of starch and related carbohydrates by randomly cleaving internal α -D-(1-4) glycosidic linkage⁸, yielding glucose, maltose, maltotriose, and other oligosaccharides⁹. It belongs to family

thirteen in the classification of glycoside hydrolases. This family is the most varied of all glycoside hydrolase families, containing many enzymes able to catalyze various reactions, such as hydrolysis, transglycosylation, condensation and cyclization¹⁰.

Amylase can be obtained from several fungi, yeast, bacteria and actinomycetes; however, enzyme from fungal and bacterial sources has dominated applications in industrial sectors¹¹. The application of an amylase in industrial reactions depends on its unique characteristics, such as its action pattern, substrate specificity, major reaction products, optimal temperature, and optimal pH¹². Bacterial α -amylase preferred for application in starch processing and textile industries due to its action at higher temperature (75-105^oC) and neutral to alkaline pH¹³. Generally production of this enzyme has been carried out by submerged fermentation^{1,2}. Among the bacterial sources *Bacillus subtilis*, *B. staerothermophilus*, *B. amyloliquefaciens*, *B. licheniformis*, *B. acidocaldarius*, *Bifidobacterium bifidum*^{1,14,15} and *B. macerans*⁵ are important species. Amylase production could be induced by starch in a stable form. When cells

were grown on maltose glucoamylase production was much lower than on starch and amylase activity disappears after 24h growth on these media¹⁶.

Designing a fermentation medium is a critical and vital process as the medium composition can significantly affect the product yield¹⁷. Screening of essential medium variables can be carried out by Plackett-Burman experimental design¹⁸. It is partial factorial design where large number of independent variables (N) are studied in small number of experiments (N+1)^{19,20}.

In the present study selected seven medium components were screened for their significance on amylase production by bacteria isolated from soil under submerged fermentation using Plackett-Burman design.

MATERIALS AND METHODS

Screening of microorganisms

Amylase producing bacteria were isolated from the soil collected from the garden of Amrutvahini College of Pharmacy, Sangamner, (Maharashtra,

India) by streak plate technique using starch agar medium. It was maintained on nutrient agar salts throughout the experiment.

Plackett-Burman experimental design

The important nutritional parameters required for enzyme production in shake flask cultures using isolated bacteria were screened from selected seven medium components *viz.* starch, yeast extract, tryptone, K₂HPO₄, CaCl₂.2H₂O, MgSO₄. 7H₂O and KCl. The Plackett-Burman experimental design of twelve trials (Table 2) for eleven variables¹⁸: seven nutrient components (independent variables) and four dummy variables (Table 1) were used to evaluate the relative importance of various nutrients. In table 2, each row represents an experiment and each column represents different variable. For each nutrient variables two different concentrations high (+) and low (-) was tested (Table 1).

Table 1: Concentrations of various variables at different levels in Plackett-Burman design for amylase production in submerged culture.

S. No.	Designation	Variables	Low level(-) g/L	High level(+) g/L
1	X ₁	Starch	5	15
2	X ₂	Yeast extract	1	5
3	X ₃	Tryptone	1	5
4	X ₄	K ₂ HPO ₄	3	7
5	X ₅	CaCl ₂ .2H ₂ O	0	0.4
6	X ₆	MgSO ₄ .7H ₂ O	0.1	0.9
7	X ₇	KCl	0	0.4
8	X ₈	Dummy1	-	-
9	X ₉	Dummy2	-	-
10	X ₁₀	Dummy3	-	-
11	X ₁₁	Dummy4	-	-

Table 2: Plackett-Burman experimental design of 12 trials for eleven variables (+) high level, (-) low level along with observed enzyme activity

Trial	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	X ₈	X ₉	X ₁₀	X ₁₁	Enzyme activity (mcg/ml)
1	+	+	-	+	+	+	-	-	-	+	-	0.423
2	-	+	+	-	+	+	+	-	-	-	+	0.245
3	+	-	+	+	-	+	+	+	-	-	-	0.284
4	-	+	-	+	+	-	+	+	+	-	-	0.261
5	-	-	+	-	+	+	-	+	+	+	-	0.164
6	-	-	-	+	-	+	+	-	+	+	+	0.125
7	+	-	-	-	+	-	+	+	-	+	+	0.437
8	+	+	-	-	-	+	-	+	+	-	+	0.409
9	+	+	+	-	-	-	+	-	+	+	-	0.384
10	-	+	+	+	-	-	-	+	-	+	+	0.157
11	+	-	+	+	+	-	-	-	+	-	+	0.336
12	-	-	-	-	-	-	-	-	-	-	-	0.144

Submerged fermentation

All experiments have been carried out in duplicates using 250 ml Erlenmeyer flasks containing 50 ml fermentation media, as per experimental design adjusted to pH 7.0 with 0.1M HCl or NaOH. These flasks were autoclaved at 15psi and 121°C for 15 minutes. Then each flask was inoculated with overnight culture of 1 ml cell suspension prepared in nutrient broth. Inoculated flasks were incubated on a rotary shaker incubator at 30°C and 150 rpm for 48 hours^{1,21}.

Enzyme extraction

Fermented medium were centrifuged to separate cells at 5°C and 5000g for 15 minutes. The clear supernants was used as crude enzyme extract²².

Enzyme assay

The activity of amylase was estimated by incubating 0.3ml of crude enzyme extract with 0.5ml 1%w/v soluble starch prepared in 0.05M phosphate buffer of pH 6.5¹⁶ for 10 min at 40°C^{23,24}. The reaction was stopped by the addition of 1ml of 3,5-dinitrosalicylic

acid reagent²⁵, the reaction mixture is heated in boiling water for 5 min. cooled and the reducing sugars released were analyzed by measuring the absorbance at 540nm⁸. An enzyme activity was expressed as the amount of reducing sugar released min⁻¹ milliliter⁻¹.

Data analysis

1. The effect of each variable was determined with the following equation²⁷:

$$E_{xi} = 2 (\sum H_{xi} - \sum L_{xi}) / N$$

where, E_{xi} is the concentration effect of the tested variable, H_{xi} and L_{xi} are the concentrations of amylase at high level and low level of the same variable, and N is the number of trials (12). When the sign is positive, the influence of variable upon enzyme production is greater at high concentration, and when the negative, the influence of variable is greater at a low concentration.

2. Mean square of each variable (the variance of effect) were calculated as follows²⁷:

$$V_{xi} = (\sum H_{xi} - \sum L_{xi})^2 / N$$

where, V_{xi} is mean square of variable.

- The experimental error was calculated by averaging the mean squares of the dummy variables²⁷:

$$R = \sum V_{xd} / n$$

where, R is the experimental error (mean square for error), V_{xd} is mean square of dummy variable, and n is number of dummy variables (2).

- Factor showing larger effects were identified using F-test²⁷:

$$F = V_{xi} / R$$

- Percentage contribution was calculated from F-test.

RESULTS

Maximum enzyme production was found in experimental trial 7, where as, minimum in 12 (Table 2) under submerged fermentation using bacteria isolated from soil. Effect of all the four dummy variables close to zero (Table 3), indicates successful experimental work. Experimental error was calculated and was found to be 0.000.

Table 3: Influence of medium variables on amylase production

Designation	Variables	ΔH	ΔL	Mean Square	Effect	F value	% Contribution
X ₁	Starch	2.273	1.096	0.115	0.196	979.89	78.06
X ₂	Yeast extract	1.879	1.490	0.013	0.065	106.89	8.51
X ₃	Tryptone	1.576	1.799	0.004	-0.038	37.02	2.95
X ₄	K ₂ HPO ₄	1.586	1.783	0.003	-0.033	27.39	2.18
X ₅	CaCl ₂ .2H ₂ O	1.866	1.503	0.011	0.061	93.28	7.43
X ₆	MgSO ₄ .7H ₂ O	1.650	1.719	0.000	-0.012	3.39	0.27
X ₇	KCl	1.736	1.633	0.001	0.017	7.47	0.60
X ₈	Dummy1	1.712	1.657	0.000	0.009	-	-
X ₉	Dummy2	1.679	1.690	0.000	-0.002	-	-
X ₁₀	Dummy3	1.690	1.679	0.000	0.002	-	-
X ₁₁	Dummy4	1.709	1.660	0.000	0.008	-	-

Among the seven selected parameters starch, yeast extract and CaCl₂.2H₂O contributes to large extent, tryptone and K₂HPO₄ have moderately significant while MgSO₄.7H₂O and KCl shows minute effect for amylase production in submerged fermentation by shake flask technique. Starch, yeast extract, CaCl₂.2H₂O and KCl were influences the production at their higher concentration; whereas, tryptone, K₂HPO₄ and MgSO₄.7H₂O were effective at lower level (fig.1).

DISCUSSION

Though amylases originate from different sources (plants, animals and microorganisms), the microbial amylases are the most produced and

used in industry, due to their productivity, thermostability, suitability over wide pH range and biocompatibility¹³. Soil and food offer the substrates for isolation of microorganism strains producing amylases. In this respect, many strains used in food industry originate from fermented food media, while soils, particularly wastes and mud offer strains used mainly in chemical industry²⁷. Production of amylase is highly dependent on starch^{1,16} hence, it have maximum contribution. Starch is ubiquitous and is an easily accessible source of energy for bacterial growth. It is composed exclusively of α - glucopyranose units that are linked to




Fig. 1: Graph showing percentage contribution and relation of factors

other by α -1,4- or α -1,6-glucosidic bonds. The two high-molecular-weight components of starch are α amylose (representing a 15 to 25% weight fraction of starch), which is a linear polymer composed exclusively of α -1,4-linked glucopyranose residues, and amylopectin (representing a 75 to 85% weight fraction of starch), which is also an α -1,4-linked glucopyranose polymer but in addition contains α -1,6-glycosidic linkages representing branch points occurring at every 17 to 26 residues. Alpha amylase is capable of amylose degradation, yielding glucose, maltose, maltotriose, and other oligosaccharides by hydrolyzing α -1,4-glucosidic bonds⁹. These reducing sugars are utilized by the bacteria for further growth as carbon source. Yeast extract was found better nitrogen

source than tryptone. Utilization of CaCl₂·2H₂O at higher level indicates metal ions Ca²⁺ also helps in enzyme activity⁸.

CONCLUSION

Designing the appropriate medium for the fermentation is an open ended, time consuming, laborious process involving large number of experiments. The Plackett-Burman experimental design is the preliminary technique for rapid illustration of the effects of various medium constituents. It tests each variable at two levels only; hence it can not give exact idea regarding the optimum level of constituent required in the medium. Therefore, further optimization of selected nutrients having greater contribution for amylase production is essential.

REFERENCES

1. Enhasy HAE, Bioprocess development for the production of α -amylase by *Bacillus amyloliquifaciens* in batch and fed-batch cultures. *Res J Microbiol* 2007; 2(7):560-568.
2. Kunamneni A, Permaul K, Singh S. Amylase production in solid state fermentation by the thermophilic fungus *Thermomyces funginosus*. *J Biosci Bioeng* 2005; 100(2):168-171.
3. Lin LL, Hsu WH, Chu WS. A gene encoding for α -amylase from thermophilic *Bacillus* sp., strain TS-23 and its expression in *Escherichia coli*. *J Appl Microbiol* 1997; 82:325-334.
4. Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D, Mohan R. Advances in microbial amylases. *Biotechnol Appl Biochem* 2000; 31:35-152
5. Aiyer PV. Amylase and their applications. *Afr J Biotechnol* 2005; 4(13):1525-1529.
6. Gupta A, Gupta VK, Modi DR, Yadava LP. Production and characterization of α -amylase from *Aspergillus niger*. *Biotechnol* 2008; 7(3):551-556.
7. Rodriguez VB, Alam eda EJ, Gallegor JF, Requena AR. Modification of the activity of an α -amylase from *Bacillus licheniformis* by several surfactants. *Electron J Biotechnol*. 2006; 9(5): DOI:10.225/vol9-issue5-fulltext-16.
8. Shaw JF, Lin FP, Chen SC, Chen HC. Purification and properties of extracellular α -amylase from *Thermus* sp., *Bot Bull Acad Sin* 1995; 36:195-200.
9. Ryan SM, Fitzgerald GF, Sinderen D. Screening for identification of starch-, amylopectin-, and pullulan-degrading activities in bifidobacterial strains. *Appl Environ Microbiol* 2006; 72(8):5289-5296.
10. Ali MB, Khemakhem B, Robert X, Haser R, Bejar S, Thermostability enhancement and change in starch hydrolysis profile of the maltohexaose-forming amylase of *Bacillus stearothermophilus* US100 strain. *Biochem J* 2006; 394:51-56.
11. Kathiresan K, Manivannan S. α -Amylase production by *Penicillium fellutanum* isolated from mangrove rhizosphere soil. *Afr J Biotechnol* 2006; 5(10):829-832.
12. Yun J, Kang S, Park S, Yoon H, Kim MJ, Heu S, Ryu S. Characterization of a novel amylolytic enzyme encoded by a gene from a soil derived metagenomic library. *Appl Microbiol Biotechnol* 2004; 70(12):7229-7235.
13. Shah NK, Ramamurthy V, Kothar RM. Comparative profile of fungal alpha amylase production by submerged and surface fermentation. *Biotech Letters* 1991; 13(5):361-364.
14. Kanno M. α -amylase production by *Bacillus acidocaldarius*, *B. stearothermophilus* and their D-cycloserine resistant mutants. *Agric Biol Chem* 1986; 50(10):2633-2635.
15. Konsoula Z, Liakopoulou-Kyrikides M. Thermostable α -amylase production by *Bacillus subtilis* entrapped in calcium alginate gel capsules. *Enz Microbiol Technol* 2006; 39(4):690-696.
16. Reyed MR. Biosynthesis and properties of extracellular amylase by encapsulation *Bifidobacterium bifidum* in batch culture. *Aust J Basic Appl Sci* 2007; 1(1):7-14.
17. Kennedy M, Krouse D. Strategies for improving fermentation medium performance: a review. *J Ind Microbiol Biotechnol* 1999; 23:456-475.
18. Plackett RL, Burman JP. The design of optimum multifactorial experiments. *Biometrika*, 1946; 33:305-325.
19. Naveena B J, Md Altaf, Bhadrariah K, Reddy G. Selection of medium component by Plackett-Burman design for production of L(+) lactic acid by *Lactobacillus amylophilus* GV6 in SSF using wheat bran. *Biores Technol* 2005; 96:485-490.
20. Sayyad SA, Panda BP, Javed S, Ali M. Screening of nutrient parameter for lovastatin production by *Monascus purpureus* MTCC 369 under submerged fermentation using Plackett-Burman design. *Res J Microbiol* 2007; 2(7):601-605.
21. Cordeiro CAM, Martins MLL, Luciano AB. Production and properties of α -amylase from thermophilic *Bacillus* sp. *Braz J Microbiol* 2002; 33(1):57-61.
22. Abu EA, Ado SA, James DB. Raw starch degrading amylase production by mixed culture of *Aspergillus niger* and *Saccharomyces cerevisiae* grown on sorghum pomace. *Afr J Biotechnol* 2005; 4(8):785-790.
23. Figueira ELZ, Hirooka EY. Culture medium for amylase production by toxigenic fungi. *Braz Arch Biol Technol* 2002; 43(5):461-467.
24. Wilson JJ, Ingledew WM. Isolation and characterization of *Schwanniomyces alluvius* amylolytic enzymes. *Appl Environ Microbiol* 1982; 44(2): 301-307.
25. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Anal Chem* 1959; 3:426-428
26. Stanbury PF, Whitkar A, Hall SJ. Principles of fermentation Technology. 2nd edition, New Delhi (India), Aditya Books (P) Ltd.; 1997.
27. Fossi BT, Tavea F, Ndjouenkeu R. Production and partial characterization of a thermostable amylase from ascomycetes yeast strain isolated from starchy soil. *Afr J Biotechnol* 2005; 4(1):14-18.